

DENDREX®

**SYSTEMIC INSECTICIDE APPLIED INTERNALLY BY TREE TECH® MICROINJECTION SYSTEMS
FOR CONTROL OF CERTAIN INSECTS ON ORNAMENTAL OR NON-BEARING TREES AND SHRUBS ONLY**

| | |
|---|--------|
| ACTIVE INGREDIENT | By Wt. |
| Acephate (O,S-dimethylacetylphosphoramidothioate) | .98% |
| OTHER INGREDIENTS | 2% |
| | 100% |

EPA Reg. No. 64014-1
EPA Est. No. 64014-FL-001

STOP - READ THE LABEL BEFORE USE

**FOR USE BY COMMERCIAL ARBORISTS
(APPLICATORS)**

KEEP OUT OF REACH OF CHILDREN

CAUTION

| FIRST AID | |
|--|--|
| Organophosphate | |
| If swallowed: | <ul style="list-style-type: none">•Call poison control center or doctor immediately for treatment advice.•Have person sip a glass of water if able to swallow.•Do not induce vomiting unless told to do so by the poison control center or doctor.•Do not give anything by mouth to an unconscious person |
| If eyes: | <ul style="list-style-type: none">•Hold eye open and rinse slowly and gently with water for 15-20 minutes.•Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.•Call a poison control center or doctor for treatment advice. |
| If on skin or clothing: | <ul style="list-style-type: none">•Take off contaminate clothing.•Rinse skin immediately with plenty of water for 15-20 minutes.•Call a poison control center or doctor for treatment advice. |
| If inhaled: | <ul style="list-style-type: none">•Move person to fresh air.•If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.•Call a poison control center or doctor for further treatment. |
| HOT LINE NUMBER | |
| Organophosphate | |
| Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-457-2022 for emergency medical treatment information. | |
| NOTE TO PHYSICIAN | |
| Acephate is a cholinesterase inhibitor. If signs of cholinesterase inhibition appear, atropine is antidotal. 2-PAM also may be used in conjunction with atropine but should not be used alone. | |

SEE REAR PANEL FOR ADDITIONAL PRECAUTIONARY STATEMENTS.

NET CONTENTS: Each microinjection unit contains 5 mL (1.5 gm a.i.) acephate systemic insecticide solution. There are 100 microinjection units (500 mL total) per case.

Sold by:
Florida Silvics, Inc.
(dba Tree Tech Microinjection Systems).
950 SE 215th Avenue
MORRISTON, FL 32668 USA
(352) 528-5335

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION



SHERRILLtree®
TREE GEAR. TREE PEOPLE.

800-525-8873

SherrillTree.com

Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, or using tobacco. Wash the outside of gloves before removing them. Remove contaminated clothing and wash clothing before reuse.

Personal Protective Equipment (PPE):

Glove that are made of any waterproof material that are chemical-resistant to this product. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Handlers must wear:

- long-sleeved shirt and long pants
- shoes plus socks
- chemical resistant gloves
- protective eyewear

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations

Users should:

1. Users should wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
2. Users should remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
3. Users should remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to birds. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Do not contaminate water when cleaning equipment or disposing of equipment wash waters.

AVISO:

PRECAUCION AL USARIO: Si Usted no puede leer o entender ingles, no use este producto hasta que la etiqueta le haya sido explicada ampliamente.

(TO THE USER: If you cannot read or understand English, do not use this product until the label has been fully explained to you.)

DIRECTIONS FOR USE

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING.

Read entire label. Use strictly in accordance with **Precautionary Statements** and **Directions**, and with applicable state and federal regulations.

Not for use on plants being grown for sale or other commercial use, or for commercial seed production, or for research purposes.

GENERAL DIRECTIONS

DENDREX is approved for use only on ornamental or non-bearing trees and shrubs. A tree or shrub is defined as non-bearing if no fruits or nuts are produced within one year of application. It is prohibited to use this product on food-bearing trees and shrubs.

Applying DENDREX using the Tree Tech microinjection system:

DENDREX label and injection instructions must be read and understood prior to use or installation of Tree Tech Microinjection Systems' microinjection units. Failure to follow these instructions may lead to injury to the installer or other persons as well as mechanical or phytotoxic damage to treated trees. The following instructions must be heeded to ensure safe and effective use of the injectors containing DENDREX insecticide.

1. Protective eye wear and rubber or neoprene gloves must be worn while handling or installing the microinjection unit to prevent accidental contact with the eyes or skin.

2. When properly installed, the microinjection unit generates internal pressure resulting in the flow of DENDREX solution through the dispenser tube. The microinjection unit must never be activated unless installed correctly and securely in the tree to be treated.

3. Microinjection units containing DENDREX may require up to several hours or more to empty depending on the health of the treated tree and local weather conditions. Never assume that microinjection units have depressurized completely because they appear nearly empty or empty. When removing injectors, individuals must wear proper eye protection and rubber or neoprene gloves. The individual should then cover the microinjection unit with one hand near the point of insertion into the stem while grasping the barrel end of the microinjection unit with the other hand. The injector should be turned slightly as it is slowly withdrawn from the tree. Careful removal of microinjection units should prevent accidental spillage of DENDREX and subsequent exposure to the installer.

4. After microinjection units are removed from treated trees they must be discarded into the heavy-duty plastic disposal bag included in



SHERRILLtree®
TREE GEAR. TREE PEOPLE.

800-525-8873

SherrillTree.com

each case of microinjection units. The bag should be properly sealed and placed in the original carton. Sealed cartons should be returned freight prepaid to Tree Tech Microinjection Systems, 950 SE 215th Ave, Morriston, FL 32668.

Installing Microinjection Units

1. Determine the number of microinjection units to be installed based upon 1) the target insect pest(s), and 2) the registered dosage rate as administered by proper spacing of microinjection units around the stem circumference. Unless otherwise noted, microinjection units should be installed in the stem and root flares every 6 inches around the trunk near the ground line, i.e., 6.0-to-8.0 inches (15-to-20 cm) from the soil surface.

2. Using a portable electric drill (600-800 rpm capacity is preferred) with a sharp, clean 11/64-in. (0.4-cm) bit, the installer should drill a hole at the correct stem circumference spacing to a depth of 1/4-to-1/2 in. (0.60-to-1.3 cm) into the wood (xylem) under the bark. A slight downwardly angle is recommended for more complete drainage of the microinjection unit.

3. After reaching the proper depth range, the drill bit should be withdrawn carefully to avoid dislodging bark fragments around the exterior opening of the hole. The rear barrel portion of the microinjection unit should be compressed partially (about 1/2 in. or 1/3 cm.) The microinjection unit should be inserted into the hole and the rear barrel portion compressed further without engaging the locking mechanism and barrel segments. Placing the plastic installation cap over the rear barrel end, strike the cap with a plastic hammer to seat the microinjection unit firmly in the hole. If the microinjection unit is not properly positioned in the hole, strike the cap again until correctly seated. By striking the microinjection unit, the back end of the feeder tip is forced back into the funnel-shaped section dislodging a septum that allows the DENDREX solution to flow from the microinjection unit into the tree.

4. When the microinjection unit is positioned correctly in the tree and the internal septum is dislodged to permit the flow of DENDREX, remove the cap and, if necessary, push the rear barrel portion of the unit further downwardly until it is flush with the edge of the Locking mechanism. This engages the locking mechanism that pressurizes the microinjection unit and assists in the evacuation of DENDREX from the microinjection unit and movement into the vascular system of the tree.

5. Each hole should be drilled and a microinjection unit installed without delay. After the unit is properly seated, it should be activated. This sequence minimizes the flow of tree sap or resin into the hole prior to DENDREX injection.

APPLICATION: Apply in all cases when plant root systems are established and actively growing. Applications should be applied at recommended intervals to maintain insect control.

| Plant | Insects | Microinjection Unit Spacing Interval Around Stem Circumference | Time of Injection |
|--|---|--|---|
| Trees (except Flowering Crabapple, see below) and Shrubs | Aphids Bagworms Birch leaf miner Tent caterpillar Lace bugs | 1 microinjection unit every 6 inches | As the insects begin to appear. |
| | Tussock moth larvae, Gypsy moth larvae | 1 microinjection unit every 6 inches | As the insects begin to appear. |
| | Scales (crawlers) | 1 microinjection unit every 6 inches | As crawlers begin to appear. Repeat applications at 4-week or more intervals. May be necessary where there is continuous crawler production. |
| | Grasshoppers | 1 microinjection unit every 6 inches | As the grasshoppers begin to appear. |
| | California oak worm Cankerworms (Spring and Fall) | 1 microinjection unit every 6 inches | As the insects begin to appear. Use the higher amount when the larger larvae are present. |
| | Nantucket pine tip moth larvae Zimmerman pine moths | 1 microinjection unit every 6 inches | Time of application is important Consult your farm advisor or County Extension Agent. Repeat applications will be required for subsequent generations. |
| | Root weevil adults | 1 microinjection unit every 6 inches | Apply when first feeding damage occurs. Repeat applications at 4-week intervals until the first heavy frost, maybe necessary for complete foliage protection. |
| | Box elder bugs Thrips White flies Sawflies Budworms Grasshoppers Cone worms Casebearer Webworms Leaf miners Pine needle miner | 1 microinjection unit every 6 inches | As the insects begin to appear. |
| | Bronze birch borer | 1 microinjection unit every 6 inches | Mid- July through August |
| | Japanese beetle | 1 microinjection unit every 4 inches | As the Japanese beetles begin to appear. Repeat applications at 4-week intervals may be necessary |
| Trees and Shrubs (continued) | Elm leaf beetle (larvae) | 1 microinjection unit every 4 inches | As the larvae begin to appear. DENDREX will not prevent elm leaf beetle eggs from hatching |
| Flowering crabapples | Aphids Tent caterpillars Leaf rollers | 1 microinjection unit every 6 inches | As the insects begin to appear |



SHERRILLtree®
TREE GEAR. TREE PEOPLE.

800-525-8873

SherrillTree.com

Caution:

Phytotoxicity has occurred on the following crabapple varieties: 'Hopa', 'Ichonoski', 'Malus floribunda', 'Pink Perfection', 'Red Wine', and 'Snow Cloud'.

GENERAL INFORMATION

STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage, or disposal of microinjection units.
Do not apply this product through any type of irrigation system.
Open dumping is prohibited.
Store microinjection units in a cool, dry place. Do not expose to temperatures below 32 F (0 C).

PESTICIDE STORAGE: Store product in the closed, original container in a cool, dry, locked place out of the reach of children.

PESTICIDE AND MICROINJECTION UNIT CONTAINER DISPOSAL: Used microinjection units may not be reused. Used microinjection units as well as any pesticide waste should be placed in the heavy-duty plastic bag that accompanies each case of microinjection units. The bag should be properly sealed, placed into the original shipping carton and returned freight prepaid for disposal to Tree Tech Microinjection Systems, 950 S.E. 215th Ave., Morriston, FL 32668.

WARRANTY AND DISCLAIMER STATEMENT

Buyer and user must follow directions when installing, removing or disposing microinjection units. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Tree Tech Microinjection Systems. The manufacturer and seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label, Tree Tech Microinjection Systems, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by the usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, buyer's or user's exclusive remedy, and Tree Tech Microinjection Systems, the manufacturer's, or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Tree Tech Microinjection Systems, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.

Label revised 01/2007

MSDS ENCLOSED